



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

certain time, the solution for this part of the motion is

$$x = \frac{Ma_1' - F}{2\rho g} \left[\cosh t \sqrt{\frac{2\rho g}{M + m' + mk^2/r^2}} - 1 \right].$$

If, at the instant $t = t_1$, a_1 changes to the constant value a_1'' , the solution for the ensuing motion is

$$x = \frac{Ma_1' - F}{2\rho g} \left[\cosh t \sqrt{\frac{2\rho g}{M + m' + mk^2/r^2}} - 1 \right] \\ + M \frac{(a_1'' - a_1')}{2\rho g} \left[\cosh (t - t_1) \sqrt{\frac{2\rho g}{M + m' + mk^2/r^2}} - 1 \right].$$

Professor Huntington's case 1 was also solved by C. C. WYLIE.

2863 [1920, 482]. Proposed by A. A. BENNETT, University of Texas.

From their generation as roulette curves, show that the two hypocycloids of five cusps drawn with common vertices, are such that each is the envelope of a chord of constant length suitably placed upon the other.

Show that for any odd prime p , the $(p - 1)/2$ distinct p -cusped hypocycloids with common vertices may be arranged in cycles, so that each is the envelope of a chord of constant length taken upon the succeeding curve of the cycle.

A solution of this problem appears on pages 371-373 of this issue of the MONTHLY.

NOTES AND NEWS.

It is to be hoped that readers of the MONTHLY will coöperate in contributing to the general interest of this department by sending items to H. P. MANNING, Brown University, Providence, R. I.

Miss GERTRUDE I. MCCAIN, of Oxford College, has been made professor of mathematics at Westminster College, New Wilmington, Pa.

Associate Professor J. V. MCKELVEY decided to remain at Iowa State College (compare 1921, 285).

We are requested to state that our note regarding Associate Professor W. A. WILSON's promotion (1921, 332) is not in accordance with fact.

At the University of Michigan, Mr. J. P. BALLANTINE, of Pennsylvania State College, and Mr. W. M. COATES, of the University of Virginia, have been appointed instructors of mathematics.

Mr. J. C. FUNK, of Tamalpais Polytechnic High School, Mill Valley, Cal., has been elected to the headship of the department of mathematics in the Santa Maria high school and junior college.

Associate Professor EMMA L. KONANTZ, who has spent two years on leave of absence teaching in Peking University, has returned to her position at Ohio Wesleyan University.

Mr. H. K. CUMMINGS, instructor of physics at the Worcester Polytechnic Institute, and experimental physicist in the research laboratories of the Acheson Graphite Company, has been appointed instructor of mathematics at Brown University.

Mr. F. W. WINTERS, of Mount Allison University, but recently a student instructor at Yale University, has been appointed assistant professor of mathematics at Dalhousie College, Halifax, Nova Scotia.